

Temporal changes of the fish community in a seagrass bed after disappearance of vegetation caused by disturbance of the sea bottom and sediment deposition

Journal:	<i>Journal of the Marine Biological Association of the United Kingdom</i>
Manuscript ID	JMBA-03-19-OA-0065.R2
Manuscript Type:	Original Article
Date Submitted by the Author:	10-Sep-2019
Complete List of Authors:	Yoshikawa, Kentaro Tomiyama, Takeshi; Graduate School of Biosphere Science, Hiroshima University Shoji, Jun
Keywords:	Natural disturbance, Seto Inland Sea, species change, succession, vegetation, <i>Zostera marina</i>
Abstract:	<p>To investigate the response of the fish community structure to a natural disturbance in their habitat, fish abundance, biomass and species composition were analyzed in relation to temporal variability of environmental conditions in a seagrass <i>Zostera marina</i> bed. A total of 3,024 fishes belonging to 46 taxa (22 families) were collected by quantitative sampling for 10 years from 2007 to 2016 in the Seto Inland Sea, southwestern Japan. Seagrass shoot density decreased to less than 1/20 of its original density after disappearance of vegetation caused by heavy rain in the fall of 2011 and the area did not recover for the next five years. In order to analyze temporal changes of fish community, the fish was divided into three groups depending on their habitats or lifestyles: pelagic or migratory species (PM), sand or mud bottom-associated species (SM) and seagrass (<i>Z. marina</i>)– or substrate (rocky bottom including macrophytes)– associated species (ZS). Multiple regression analysis showed seagrass shoot density had the most significant effect on biomass of ZS among the three groups, with higher fish biomass under higher seagrass shoot density. Fish community composition changed after the disappearance of the seagrass vegetation coverage with an increase in abundance of SM during the five-years of the post-disturbance period. Seagrass vegetation was concluded to affect temporal change of fish community structure through a stronger influence on fish species that are more dependent on seagrass bed as habitat.</p>

SCHOLARONE™
Manuscripts

1

Running Head:

2

Temporal change of fish community in seagrass bed

3

4

5

Title:

6

Temporal changes of the fish community in a seagrass bed after

7

disappearance of vegetation caused by disturbance of the sea

8

bottom and sediment deposition

9

10

11

KENTARO YOSHIKAWA¹, TAKESHI TOMIYAMA¹ and JUN SHOJI^{1, 2}

12

13

14

15

¹*Graduate School of Biosphere Science, Hiroshima University, 1-4-4 Kagamiyama, Higashi-*

16

hiroshima, Hiroshima 739-8528, Japan

17

18

²*Present address: Atmosphere and Ocean Research Institute, The University of Tokyo, 5-1-5*

19

Kashiwanoha, Kashiwa, Chiba 277-8564, Japan

20

21

22

23

Corresponding author:

24

K. Yoshikawa

25

E-mail: m182767@hiroshima-u.ac.jp

26

Abstract

To investigate the response of the fish community structure to a natural disturbance in their habitat, fish abundance, biomass and species composition were analyzed in relation to temporal variability of environmental conditions in a seagrass *Zostera marina* bed. A total of 3,024 fishes belonging to 46 taxa (22 families) were collected by quantitative sampling for 10 years from 2007 to 2016 in the Seto Inland Sea, southwestern Japan. Seagrass shoot density decreased to less than 1/20 of its original density after disappearance of vegetation caused by heavy rain in the fall of 2011 and the area did not recover for the next five years. In order to analyze temporal changes of fish community, the fish was divided into three groups depending on their habitats or lifestyles: pelagic or migratory species (PM), sand or mud bottom-associated species (SM) and seagrass (*Z. marina*)- or substrate (rocky bottom including macrophytes)- associated species (ZS). Multiple regression analysis showed seagrass shoot density had the most significant effect on biomass of ZS among the three groups, with higher fish biomass under higher seagrass shoot density. Fish community composition changed after the disappearance of the seagrass vegetation coverage with an increase in abundance of SM during the five-years of the post-disturbance period. Seagrass vegetation was concluded to affect temporal change of fish community structure through a stronger influence on fish species that are more dependent on seagrass bed as habitat.

Keywords: Natural disturbance; Seto Inland Sea; species change; succession; vegetation; *Zostera marina*

INTRODUCTION

Disturbance caused by anthropogenic and natural forces on an ecosystem is known to greatly affect the plant and animal communities (Foster *et al.*, 1998; Nyström *et al.*, 2000; Johnson & Miyanishi, 2010; Turner, 2010). Some of the disturbance caused by natural forces happens suddenly and is therefore difficult to predict. To evaluate the effects of a natural disturbance, it is necessary to collect and compare data before and after the event. In addition to the data collection just before and after the event, long-term monitoring as well as spatial comparison would also be useful to clarify the process of community succession (Larkum *et al.*, 2007; Jelbart *et al.*, 2007; Hori *et al.*, 2009).

In marine coastal ecosystems, flooding and tsunamis are typical examples of natural disturbance that directly affect ecosystems and plant and animal communities owing to organisms being swept

away and to the drastic changes in e.g. physical, chemical and biological properties of their habitats (Atwater & Moore 1992; Nakaoka *et al.*, 2006; Jaramillo *et al.*, 2012). Furthermore, biological communities at higher trophic levels suffer from indirect effects through changes in the habitat conditions (e.g. bottom sediments and plant vegetation: Muraoka *et al.* 2017). However, information on the effect of natural disturbances on the fish community is very limited (Shoji & Morimoto, 2016; Noda *et al.*, 2017) while there have been observations on the short-term impact of the disturbances on the benthic communities on seagrass beds (Nakaoka *et al.*, 2006; Whanpetch *et al.*, 2010), the sandy bottom (Seike *et al.*, 2013) and rocky reefs (Jaramillo *et al.*, 2012; Takami *et al.*, 2013). The difficulty in quantitative sampling of fish and the greater effort required for fish surveys have restricted the evaluation of the effects of natural disturbance on fish communities (Beck *et al.*, 2000). Monitoring of the fish community through the periods before and after a natural disturbance is indispensable to better understand the impacts of disturbance.

Seagrass *Zostera marina* beds are one of the most important ecosystems that serve as a habitat for a variety of marine organisms. Among the world's ecosystems, seagrass (including *Zostera* spp. and related species) beds provide ecosystem services with high economic values (Costanza, 1997; Ellison *et al.*, 2005; Orth *et al.*, 2006). Many fish species are dependent on seagrass beds for their spawning, feeding, refuge from predators and inhabitation (Heck *et al.*, 1989; Boström *et al.*, 2006; Larkum *et al.*, 2007). The abundance of fishes associated with seagrass beds has been reported to fluctuate depending on spatial and temporal variabilities in seagrass vegetation (Jelbart *et al.*, 2007; Hori *et al.*, 2009; Raventos *et al.*, 2009; Shoji *et al.*, 2017). Changes in fish community after the decline of seagrass beds resulting from eutrophication, physical disturbance and a resurgence of wasting disease have been observed in the North Atlantic (Hughes *et al.*, 2002). In recent studies, drastic changes in the fish community in seagrass beds following an abrupt decrease in seagrass vegetation coverage have been reported. On the Pacific coast of northern Japan after the tsunami following the 2011 Tohoku earthquake off the Pacific coast, the dominant fish species (fishes associated with seagrass beds and substrates: rocky bottom including macrophytes) changed to sand or mud bottom-associated fish species after the disturbance of bottom sediment (Shoji & Morimoto, 2016; Noda *et al.*, 2017). In these previous studies, however, the periods of the observations (one to three years) are limited just before and after the natural disturbances caused by drastic changes in the seagrass vegetation and sea bottom. There is no study that analyzed the effect of drastic changes of seagrass vegetation coverage on the fish community based on monitoring for longer periods.

In the present study, data on environmental conditions and fish community in a seagrass bed in the Seto Inland Sea, Japan, affected by a disturbance of the sea bottom and sediment deposition following heavy rain in 2011 was analyzed from 2007 to 2016. The hypothesis that temporal variability in

seagrass vegetation coverage induces change of fish community structure through a more significant effect on seagrass or substrate-associated fish species was tested.

MATERIALS AND METHODS

Field survey

Surveys for seagrass vegetation and the fish community were conducted on a seagrass bed (ca. 10 ha) off the eastern Ikuno Island, central Seto Inland Sea, Japan (34°17'20"N, 132°55'32"E; Figure 1). Ikuno Island has a population of approximately 17, with no human habitation on the eastern coast. The vegetation of the seagrass bed is dominated by the seagrass *Z. marina*, and the mean shoot density of this plant around the sampling site fluctuates between 20 and 160 m⁻² throughout the year (Mohri *et al.*, 2013). The bottom of the seagrass area is composed of mud and sand. A heavy rain in the fall of 2011 induced disturbance of the sea bottom and sediment deposition on the seagrass bed on the eastern shore of Ikuno Island and the area did not recover for the next five years (see the Results).

Fish sampling was conducted using a round seine net (2 m high, 30 m long, and 4 mm mesh aperture; Kamimura & Shoji, 2013) in the day (1100–1700 h) during the spring tide period in August or September from 2007 to 2016. Fish were collected from four separate locations randomly selected from areas within the seagrass bed (four replicates). Tidal levels were between 50–130 cm (within two hours before and after low tide), when the shore line was close to the edge of the seagrass bed. Three sides of a square (10 m in side length) were surrounded using the net at a speed of ca. 1.0 m/s, with another side facing into the shore (around the border of the seagrass bed). Then the net was pulled landward. Each fish collection covered an area of 100 m². The collected fish were preserved in 10% formalin seawater solution. The temperature and salinity of the surface water were measured at each sampling. Seagrass shoot density was measured in at least four randomly placed 0.5 m square quadrats in the seagrass bed. The length of seagrass leaves from at least 10 shoots was measured.

Data analysis

In the laboratory, fish were identified according to Nakabo (2013). Mean number of fish species (no. fish species 100 m⁻²), abundance (no. fish 100 m⁻²) and biomass (wet weight of fish 100 m⁻²) were calculated. The total length (TL, mm) of each fish was measured to the nearest 0.1 mm. To detect the possible effect of the disturbance on seagrass vegetation and the fish community, the mean seagrass shoot density and leaf length, number of species, abundance, and biomass were compared between the periods before and after the event in 2011 using the Mann-Whitney *U* test.

In the previous studies conducted in northern Japan, the effects of the temporal change in seagrass

vegetation on fish species that were associated with the seagrass bed and substrates (rocky bottom including macrophytes) were suggested to be more intensive compared to other fish species (Shoji & Morimoto, 2016; Noda *et al.*, 2017). The collected fishes were divided into three groups according to these previous studies: pelagic or migratory species (PM), sand or mud bottom-associated species (SM) and seagrass (*Z. marina*) or substrate (rocky bottom including macrophytes)-associated species (ZS). In order to examine possible effects of the environmental conditions on the fish community, a linear model was constructed with the mean seagrass shoot density, water temperature and salinity as explanatory variables and mean biomass of the three fish groups as response variables. The leaf length was not included in the analysis due to positive correlation with seagrass shoot density ($r^2=0.680$, $p<0.01$). The model selection was operated based on the Akaike information criterion. Data of *Sebastes* spp. (juveniles) in 2012 was excluded from the analysis because of a significant effect of its dominance (90.3% in biomass: see the Results). The Bray-Curtis dissimilarity index was calculated for each year based on fish abundance and was processed for nonmetric multidimensional scaling (NMDS) to visualize the differences in the fish community for each year because the index has been applied for comparison of marine animal community structures (Clarke, 1993; Field *et al.*, 1982; Warwick & Clarke, 2001). All statistical analyses were performed in R (3.4.0: R Development Core Team).

Results

Physical environmental conditions and seagrass vegetation

The water temperature ranged between 24.7 °C (2013) and 27.8 °C (2010) and salinity ranged between 27.8 (2013) and 33.0 (2015 and 2016: Table 1). The mean (\pm standard deviation: SD) seagrass shoot density ranged between 0.3 ± 0.5 shoots m^{-2} (2015) and 61.8 ± 20.2 shoots m^{-2} (2007) (Figure 2a). Difference in the mean seagrass shoot density between the periods before and after the disturbance was significant (Mann-Whitney *U* test, $p<0.01$).

Fish community

A total of 3,024 fishes belonging to 46 taxa (22 families) were collected during the 10-year survey (Table 1). The mean (\pm SD) number of fish species ranged between 4.75 ± 0.5 100 m^{-2} (2014) and 12.5 ± 2.9 100 m^{-2} (2011: Figure 2b). The mean number of fish species decreased after the disturbance and was the lowest in 2014, with a significant difference between the two periods (Mann-Whitney *U* test, $p<0.05$).

The mean (\pm SD) fish abundance per 100 m^2 ranged between 17.0 ± 9.3 (2014) and 220.0 ± 250.2 (2012: Figure 2c). The mean (\pm SD) fish biomass per 100 m^2 ranged between 18.5 ± 2.6 g (2014) and 504.5 ± 556.1 g (2012: Figure 2d). In 2012, one year post-disturbance, the mean fish abundance,

biomass and their SDs increased due to the presence of juvenile black rockfish *Sebastes* spp. at a remarkably high level of abundance in one of the four collection areas within a scattered seagrass bed (Table 1). The differences in fish abundance and biomass between the two periods were not significant (Mann-Whitney U test, $p=0.42$ for abundance and $p=0.31$ for biomass).

Among the three fish groups (PM, SM and ZS, divided based on their habitats and lifestyles), year-to-year variability in the biomass were larger in SM and ZS (Figure 3). The seagrass shoot density and salinity were selected as explanatory variables for ZS in the model, with higher fish biomass under higher seagrass shoot density and salinity ($r^2=0.75$: Table 2). For ZS, the effect of seagrass shoot density was significant ($p=0.002$) and much greater than that of salinity ($p=0.077$). All initial explanatory variables were not significant ($p>0.6$) and were excluded by the model for SM. For PM type fish, all initial explanatory variables were adopted in the selected model ($r^2=0.28$) but were not significant ($p>0.1$). Based on the fish community structures, the years were divided into three or more groups (stress=0.132, Figure 4). Group 1: 2007 and 2011; group 2: 2008, 2009, 2010 and 2012; group 3: 2014 and 2015 and outgroup: 2013 and 2016.

DISCUSSION

Effects of environmental conditions on the fish community

The area of seagrass bed has been decreasing at a rate of 5% per year or more on the entire earth (Waycott *et al.* 2009). The effects of human activity (e.g. low oxygenation concentrations and high turbidity due to eutrophication) and natural effects are considered to affect the seagrass growth and vegetation (Larkum *et al.*, 2007). As global warming progresses, fish would also be indirectly affected through the effects on seagrass. Because the seagrass is vulnerable to high water temperature, the area of the seagrass, which is important as a fish habitat, can decrease under the global warming (Kuwaie & Hori, 2019). In addition, the global warming tends to increase the frequency of heavy rains and amount of single rainfall, so that can also cause disturbance to the bottom sediments of seagrass bed. Mitigating the effects of human activity will reduce the rate of disappearance of the seagrass beds. Furthermore, it is possible to prepare for improvement of recovery from environmental fluctuation and damage in the future through maintenance of population structure/network which enables to supply seagrass seeds from the surrounding area when seagrass beds disappear (Larkum *et al.*, 2007; Kuwaie & Hori, 2019).

In the present study, seagrass shoot density showed the significant effect only on seagrass- or substrate-associated (ZS) fish species among the three fish groups (Table 2). In general, spatial and temporal variations in habitat complexity and connectivity to adjacent habitats affect fish community

structures of seagrass beds and surrounding areas (Dorenbosch *et al.*, 2005; Dorenbosch *et al.*, 2006; Grol *et al.*, 2011). Previous studies have shown higher number of fish species, and greater abundance and biomass of fishes in areas with seagrass vegetation coverage, compared to those in surrounding areas with less or without seagrass vegetation (Ferrel & Bell, 1991; Larkum *et al.*, 2007). It is likely that the three-dimensional habitat complexity provided by seagrass leaves serves as predation refuge, feeding ground and as a habitat for fishes. In the present study, it was demonstrated that year-to-year variability of seagrass shoot density was one of the important determinants for the ZS-type fish.

Salinity was the important factor for ZS- and PM-type fish species although the effect was not significant. Seagrass is widely distributed in estuarine waters with high tolerance to low salinity condition (Nakaoka & Aioi, 2001). In a laboratory experiment conducted at salinities between 0-33 (0, 5, 10, 15, 20, 25 and 33), seagrass showed the highest germination rates at salinity of 0 under five temperatures tested (5, 10, 15, 20 and 25°C: Yamaki *et al.*, 2006). Therefore, it is plausible that the temporal variability in salinity (especially, low salinity conditions) caused by the heavy rain did not have negative effect on the seagrass vegetation in the present study site. The indirect effect of salinity through seagrass vegetation on fish community and direct effect of salinity on fish are also suggested to be minimal because the variability of salinity recorded in the seagrass bed of the study site (27-33) was relatively small (Nakaoka & Aioi, 2001; Larkum *et al.*, 2007).

Temporal variability of seagrass vegetation

Flora and fauna in coastal habitats were generally subject to high variability in environmental conditions such as temperature, salinity, dissolved oxygen concentration and turbidity at a variety of temporal scales affected by tide, freshwater flow and anthropogenic effects (Schubel, 1968; Valiela *et al.*, 1992). The succession process of the plant vegetation and the animals after a strong disturbance in marine ecosystems seems to differ based on the level of disturbance. In seagrass beds, there have been studies on the effects of natural disturbance such as disturbance of sea bottom and sediment deposition on seagrass vegetation (Nakaoka *et al.*, 2006), benthic macrofauna (Whanpetch *et al.*, 2010) and fishes (Shoji & Morimoto, 2016; Noda *et al.*, 2017). In the present survey area, the seagrass vegetation drastically decreased in 2012 (46.0 shoots m⁻² in 2011 to 8.0 shoots m⁻² in 2012). The seagrass shoot density has remained lower in recent years than that in the years before 2011. So far, the seagrass vegetation coverage has not increased in the present survey site, although there are plenty of seagrass beds with high shoot density in the surrounding waters. These seagrass beds in the surrounding waters have not experienced a loss of vegetation coverage in recent years and have most likely been able to provide the present survey site with seagrass seeds. Therefore, there may be other continuous factors that have been preventing seagrass growth in the present survey site. Turbidity in the seagrass bed has

increased since the inflow of mud caused by the heavy rain in the autumn of 2011. Additionally, an increase in the abundance of herbivore fishes such as *Siganus fuscescens* (Table 1) may have potentially affected seagrass growth and coverage.

Temporal change in dominant fish species

The dominant fish species were replaced in the present survey area after the decrease of vegetation in 2011. The ZS-type fishes such as *Sebastes oblongue*, *Hypodytes rubripinnis* and *Pterogobius elapoides* had been continuously collected in the seagrass bed before 2011 and were not collected in most of the years after 2011. The decreases in the total fish species richness, abundance and biomass (except for 2012) after 2011 indicate the loss of habitat provided by seagrass vegetation, which affected ZS-type fishes the most. The loss and decrease in habitat and its complexity also can alter the growth and survival of young fishes by affecting feeding conditions and the seagrass' function as a predation refuge (Larkum *et al.*, 2007). Therefore, temporal changes in vegetation coverage have a high potential for impact on fish species richness, abundance, and biomass in seagrass beds even within the same location. On the other hand, SM-type fishes such as *Pagrus major*, *Sillago japonica* and *Repomucenus beniteguri* were more frequently collected in the seagrass bed after 2012. The seagrass beds surveyed in the present study with decreased vegetation are suggested more suitable for these fish species that are associated with sand or mud bottom after 2012.

In 2012, juveniles of *Sebastes* spp. were collected at a high mean abundance (207.8 100 m⁻²). These *Sebastes* juveniles migrate into seagrass and macroalgal beds at about 20 mm in total length in the central Seto Inland Sea (Kamimura & Shoji, 2009). In a previous study, occurrence of three *Sebastes* juveniles (*Sebastes inermis*, *S. ventricosus* and *S. cheni*) was reported in this area (Kamimura *et al.*, 2013). Among the three species, juvenile *S. cheni* was most dominant accounting for 77.6 and 80.0% in number of the three species in 2007 and 2008, respectively (Kamimura *et al.*, 2011). The mean abundance of *S. cheni* juvenile in 2008 (451.2 individuals 100 m⁻²) was higher than that observed in the present study in 2012 (207.8 individuals 100 m⁻²), indicating a large inter-annual variability of juvenile recruits. In 2012, aggregation of the *Sebastes* juveniles around the scattered seagrass bed after recruiting at a high abundance might have induced the high mean juvenile abundance although the seagrass shoot density was low.

In summary, the fish community in the seagrass bed off Ikuno Island was dominated by seagrass- or substrate-associated species during the pre-disturbance period with high vegetation coverage. After the disappearance of vegetation caused by the heavy rain of 2011, fish species richness, abundance and biomass decreased due to the absence of the species belonging to these dominant fish group. During the five years after the heavy rain, there was no significant recovery of seagrass vegetation

268 coverage. The species richness increased to the same level as that before the heavy rain due to the
269 increase in that of sand or mud bottom-associated species. The seagrass- or substrate-associated
270 species, that were dominant before the seagrass vegetation loss, were replaced with sand or mud
271 bottom-associated species.

272

273

274 **ACKNOWLEDGEMENTS**

275

276 We express our thanks to Mr. Sadaharu Iwasaki and the staff of Takehara Marine Science Station,
277 Hiroshima University, for their assistance in the field survey. Thanks to two anonymous reviewers for
278 their constructive comments on the earlier version of the manuscript.

279

280 **FINANCIAL SUPPORT**

281

282 This study was partially supported by the Global Environment Research Fund (J.S., grant number RF-
283 0907and RF-1102) of the Ministry of the Environment, and Grant-in-Aid for Scientific Research (B)
284 (J.S., grant number 24380107).

285

286 **REFERENCES**

287

288 **Atwater BF and Moore AL** (1992) A tsunami about 1000 years ago in Puget Sound, Washington.
289 *Science* **258**, 1614–1617.

290 **Beck MW, Heck KL, Able KW, Childers DL, Eggleston DB, Gillanders BM, Halpern B, Hays**
291 **CG, Hoshino K, Minello TJ, Orth RJ, Sheridan PF and Weinstein MP** (2001) The identification,
292 conservation, and management of estuarine and marine nurseries for fish and invertebrates.
293 *Bioscience* **51**, 633–641.

294 **Boström C, Jackson EL and Simenstad CA** (2006) Seagrass landscapes and their effects on
295 associated fauna: a review. *Estuarine, Coastal and Shelf Science* **68**, 383–403.

296 **Clarke KR** (1993) Non - parametric multivariate analyses of changes in community structure.
297 *Australian journal of ecology* **18**: 117–143.

298 **Costanza R, d’Arge R, de Groot R, Farber S, Grasso M, Hannon B, Limburg K, Naeem S,**
299 **O’Neill RV, Paruelo J, Raskin RG, Sutton P and Van den Belt M** (1997) The value of the world’s
300 ecosystem services and natural capital. *Nature* **387**, 253–260.

- 301 **Dorenbosch M, Grol MGG, Nagelkerken I, and Van der Velde G** (2006). Different surrounding
302 landscapes may result in different fish assemblages in East African seagrass beds. *Hydrobiologia*,
303 **563**, 45–60.
- 304 **Dorenbosch M, Grol MGG, Christianen MJA, Nagelkerken I, and Van Der Velde G** (2005). Indo-
305 Pacific seagrass beds and mangroves contribute to fish density and diversity on adjacent coral reefs.
306 *Marine Ecology Progress Series*, **302**, 63–76.
- 307 **Ellison AM, Bank MS, Clinton BD, Colburn EA, Elliott K, Ford CR and Mohan J** (2005) Loss
308 of foundation species: consequences for the structure and dynamics of forested
309 ecosystems. *Frontiers in Ecology and the Environment* **3**, 479–486.
- 310 **Ferrel DJ and Bell JD** (1991) Differences among assemblages of fish associated with *Zostera*
311 *capricorni* and bare sand over a large spatial scale. *Marine Ecology Progress Series* **72**, 15–24.
- 312 **Field JG, Clarke KR and Warwick RM** (1982) A practical strategy for analysing mutispecies
313 distribution patterns. *Marine Ecology Progress Series* **8**, 37–52.
- 314 **Foster DR, Knight DH and Franklin JF** (1998) Landscape patterns and legacies resulting from large,
315 infrequent forest disturbances. *Ecosystems* **1**, 497–510.
- 316 **Grol MG, Nagelkerken I, Rypel AL, and Layman CA** (2011) Simple ecological trade-offs give rise
317 to emergent cross-ecosystem distributions of a coral reef fish. *Oecologia* **165**, 79–88.
- 318 **Heck KL, Able KW, Fahay MP and Roman CT** (1989) Fishes and decapod crustaceans of Cape
319 Cod eelgrass meadows: species composition, seasonal abundance patterns and comparison with
320 unvegetated substrates. *Estuaries* **12**, 59–65.
- 321 **Hori M, Suzuki T, Monthum Y, Srisombat T, Tanaka Y, Nakaoka M and Mukai H** (2009) High
322 seagrass diversity and canopy-height increase associated fish diversity and abundance. *Marine*
323 *Biology* **156**, 1447–1458.
- 324 **Hughes JE, Deegan LA, Wyda JC, Weaver MJ and Wright A** (2002) The effects of eelgrass habitat
325 loss on estuarine fish communities of southern New England. *Estuaries* **25**, 235–249.
- 326 **Jaramillo E, Dugan JE, Hubbard DM, Melnick D, Manzano M, Duarte C, Campos C and**
327 **Sanchez R** (2012) Ecological implications of extreme events: footprints of the 2010 earthquake
328 along the Chilean coast. *PLoS ONE* **7**, e35348. doi: 10.1371/journal.pone.0035348.
- 329 **Jelbart JE, Ross PM and Connolly RM** (2007) Patterns of small fish distributions in seagrass beds
330 in a temperate Australian estuary. *Journal of the Marine Biological Association of the United*
331 *Kingdom* **87**, 1297–1307.
- 332 **Johnson EA and Miyanishi K** (2010) *Plant disturbance ecology: the process and the response*.
333 Amsterdam: Elsevier.

- 334 **Kamimura Y and Shoji J** (2009) Seasonal changes in the fish assemblage in a mixed vegetation area
 335 of seagrass and macroalgae in the central Seto Inland Sea. *Aquaculture Science* **57**, 233-241.
- 336 **Kamimura Y and Shoji J** (2013) Effects of vegetation coverage on post-settlement growth, survival
 337 and recruitment potential of juvenile black rockfish *Sebastes cheni* in a macroalgal bed. *Estuarine,*
 338 *Coastal and Shelf Science* **129**, 86–93.
- 339 **Kamimura Y, Kasai A and Shoji J** (2011) Production and prey source of juvenile black rockfish
 340 *Sebastes cheni* in a seagrass and macroalgal bed in the Seto Inland Sea, Japan: estimation of the
 341 economic value of a nursery. *Aquatic Ecology* **45**, 367–376.
- 342 **Kamimura Y, Kawane M, Hamaguchi M and Shoji J** (2013) Age and growth of three rockfish
 343 species, *Sebastes inermis*, *S. ventricosus* and *S. cheni*, in the central Seto Inland Sea, Japan.
 344 *Ichthyological Research* **61**, 108-114.
- 345 **Kuwae T and Hori M** (2019) *Blue Carbon in Shallow Coastal Ecosystems: Carbon Dynamics, Policy,*
 346 *and Implementation*. Springer.
- 347 **Larkum AWD, Orth RJ and Duarte CM** (2007) *Seagrasses: Biology, Ecology and Conservation*.
 348 Dordrecht: Springer.
- 349 **Mohri K, Kamimura Y, Mizuno K, Kinoshita H, Toshito S and Shoji J** (2013) Seasonal changes
 350 in the fish assemblage in a seagrass bed in the central Seto Inland Sea. *Aquaculture Science* **61**, 215-
 351 220.
- 352 **Muraoka D, Tamaki H, Takami H, Kurita Y and Kawamura T** (2017) Effects of the 2011 Great
 353 East Japan Earthquake and tsunami on two kelp bed communities on the Sanriku coast. *Fisheries*
 354 *Oceanography* **22**, 128–140.
- 355 **Nakabo T** (2013) *Fishes of Japan: with pictorial keys to the species*. 3rd edition. Tokyo: Tokai
 356 University Press.
- 357 **Nakaoka M and Aioi K** (2001) Ecology of seagrasses *Zostera* spp. (Zosteraceae) in Japanese waters:
 358 a review. *Otsuchi Marine Science* **26**, 7–22.
- 359 **Nakaoka M, Tanaka Y, Mukai H, Suzuki T and Aryuthata C** (2006) Tsunami impacts on
 360 biodiversity of seagrass communities in the Andaman Sea, Thailand: (1) Seagrass abundance and
 361 diversity. *The Nagisa World Congress* **8**, 49–56.
- 362 **Noda T, Hamaguchi H, Fujinami Y, Shimizu D, Aono H, Nagakura Y, Fukuta A, Nakano H,**
 363 **Kamimura Y and Shoji J** (2017) Impact of the tsunami caused by the Great East Japan Earthquake
 364 on seagrass beds and fish communities in Miyako Bay, Japan. *Coastal Ecosystems* **4**, 12–25.
- 365 **Nyström M, Folke C and Moberg F** (2000) Coral reef disturbance and resilience in a human-
 366 dominated environment. *Trends in Ecology & Evolution* **15**, 413–417.

- 367 **Orth RJ, Carruthers TJB, Dennison WC, Duarte CM, Fourqurean JW, Heck KL, Huches Jr.**
368 **AR, Kendrick GA, Kenworthy WJ, Olyarnik S, Short FT, Waycott M and Williams SL** (2006)
369 A global crisis for seagrass ecosystems. *Bioscience* **56**, 987–996
- 370 **Raventos N, Ferrari B and Planes S** (2009) Differences in population parameters and behaviour of
371 the herbivorous fish *Sarpa salpa* between protected and unprotected seagrass meadows in the north-
372 western Mediterranean. *Journal of the Marine Biological Association of the United Kingdom* **89**,
373 1153–1159.
- 374 **Schubel JR** (1968) Turbidity maximum of the northern Chesapeake Bay. *Science* **161**, 1013e1015.
- 375 **Seike K, Shirai K and Kogure Y** (2013) Disturbance of shallow marine soft-bottom environments
376 and megabenthos assemblages by a huge tsunami induced by the 2011 M9.0 Tohoku-Oki
377 earthquake. *PLoS One* **8**, e65417. doi: 10.1371/journal.pone.0065417.
- 378 **Shoji J and Morimoto M** (2016) Changes in fish community in seagrass beds in Mangoku-ura Bay
379 from 2009 to 2014, the period before and after the tsunami following the 2011 off the Pacific coast
380 of Tohoku earthquake. *Journal of Oceanography* **72**, 91–98.
- 381 **Shoji J, Mitamura H, Ichikawa K, Kinoshita H and Arai N** (2017). Increase in predation risk and
382 trophic level induced by nocturnal visits of piscivorous fishes in a temperate seagrass bed. *Scientific*
383 *Reports* **7**, 3895.
- 384 **Takami H, Won NI and Kawamura T** (2013) Impacts of the 2011 mega-earthquake and tsunami on
385 abalone *Haliotis discus hannai* and sea urchin *Strongylocentrotus nudus* populations at Oshika
386 Peninsula, Miyagi, Japan. *Fisheries Oceanography* **22**, 113–120.
- 387 **Turner M** (2010) Disturbance and landscape dynamics in a changing world. *Ecology* **91**, 2833–2849.
- 388 **Valiela I, Foreman K, LaMontagne M, Hersh D, Costa J, Peckol P, DeMeo-Anderson B,**
389 **D’Avanzo C, Babione M, Sham CH, Brawley J and Lajtha K** (1992) Couplings of watersheds
390 and coastal waters: sources and consequences of nutrient enrichment in Waquoit Bay,
391 Massachusetts. *Estuaries* **15**, 443–457.
- 392 **Warwick RM and Clarke KR** (2001) Practical measures of marine biodiversity based on relatedness
393 of species. *Oceanography and Marine Biology: An Annual Review* **39**, 207–231.
- 394 **Waycott M, Duarte CM, Carruthers TJB, Orth RJ, Dennison WC, Olyarnik S, Calladine A,**
395 **Fourqurean JW, Heck Jr. KL, Hughes AR, Kendrick GA, Kenworthy WJ, Short FT and**
396 **Williams SL** (2009) Accelerating loss of seagrasses across the globe threatens coastal ecosystems.
397 *Proceedings of the National Academy of Sciences* **106**, 12377–12381.

398 **Whanpetch N, Nakaoka M, Mukai H, Suzuki T, Nojima S, Kawai T and Aryuthaka C (2011)**
399 Temporal changes in benthic communities of seagrass beds impacted by a tsunami in the Andaman
400 Sea, Thailand. *Estuarine, Coastal and Shelf Science* **87**, 246–252.

401 **Yamaki K, Ogawa H, Yoshikawa H and Nanba N (2006)** Effects of salinity and temperature on the
402 germination activation of eelgrass seeds. *Aquaculture Science* **54**, 347–351.

403

404 **FIGURE CAPTIONS**

405

406 **Fig. 1.** Map showing the location of the seagrass bed off Ikuno Island, Hiroshima Prefecture, western
407 Japan, where environmental survey and fish collection were conducted from 2007 to 2016. Depth
408 contours of 10 and 20 m are indicated as dotted lines.

409

410 **Fig. 2.** Mean shoot density of seagrass (number of shoots 100 m⁻²: a), mean number of fish species (b),
411 fish abundance (number of fish 100 m⁻²; c) and fish biomass (g 100 m⁻²; d) collected in the seagrass
412 bed off Ikuno Island from 2007 to 2016. Dotted lines and the vertical bars indicate the disturbance by
413 the heavy rain in fall 2011 and standard deviation, respectively. Photographs on top of the figure show
414 an underwater overview of the seagrass bed in 2008 and 2014.


415

416 **Fig. 3.** Mean biomass (wet weight 100 m⁻²) of three fish groups based on their habitat and/or life cycles
417 from 2007 to 2016. PM: pelagic and migrative group, SM: sand- or mud-bottom associated group, ZS:
418 seagrass- or substrate-associated group.

419

420 **Fig. 4.** Non-metric multidimensional scaling (NMDS) ordination using the Bray-Curtis dissimilarity
421 index to differentiate the fish species composition in the seagrass bed off Ikuno Island from 2007 to
422 2016.

423







Fig. 2. Mean shoot density of seagrass (number of shoots 100 ⁻²: a), mean number of fish species (b), fish abundance (number of fish 100 m⁻²; c) and fish biomass (g 100 m⁻²; d) collected in the seagrass bed off Ikuno Island from 2007 to 2016. Dotted lines and the vertical bars indicate the disturbance by the heavy rain in fall 2011 and standard deviation, respectively. Photographs on top of the figure show an underwater overview of the seagrass bed in 2008 and 2014.

111x185mm (72 x 72 DPI)






Table 1. Mean fish abundance (number 100 m⁻²), biomass (wet weight 100 m⁻²), water temperature (°C), and salinity at the seagrass bed off Ikuno Island, Seto Inland Sea, western Japan, from 2007 to 2016. Fish were divided into 3 groups driven from each habitat and/or lifestyles, PM: pelagic or migratory species, SM: sand or mud bottom-associated species, ZS: seagrass (*Z. marina*)- or substrate (rocky bottom including macrophytes)-associated species.

Family	Species	Group	No. of ind. / 100 m ²												Wet weight (g / 100 m ²)											
			2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Total	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	Total		
Engraulidae	<i>Engraulis japonica</i>	PM					5.0					0.5	5.5					2.4					0.4	2.7		
Syngnathidae	<i>Urocampus nanus</i>	ZS					0.3						0.3					0.03						0.0		
	<i>Syngnathus schlegeli</i>	ZS		1.0	0.8		0.5	0.5	0.5	1.3	1.5	6.0		1.4	1.4		0.1	0.1		0.2	0.7	1.6	5.5			
	<i>Hippocampus coronatus</i>	ZS				1.5						0.3	1.8			0.5						0.1	0.6			
Mugilidae	<i>Mugil cephalus cephalus</i>	PM		2.0			3.0		0.5				5.5		2.5		6.0		0.4				8.9			
	<i>Chelon haematocheilus</i>	PM	7.8		0.8	0.3							8.8	14.5		1.2	0.5						16.2			
Hemiramphidae	<i>Hyporhamphus sajori</i>	PM	0.3										0.3	1.2									1.2			
Sebastidae	<i>Sebastes schlegelii</i>	ZS					0.8						0.8				3.5						3.5			
	<i>Sebastes oblongus</i>	ZS	0.8	0.5	1.0	0.3							2.5	2.3	1.5	3.6	0.9						8.2			
	<i>Sebastes</i> spp.	ZS	36.8	41.3	11.0	14.3	3.0	207.8	0.8			12.0	326.8	205.8	166.6	43.8	136.9	15.5	471.3	3.7			33.8	1077.3		
Tetrarogidae	<i>Hypodytes rubripinnis</i>	ZS	2.0	0.8	0.8	0.5	0.3	0.3					4.5	20.6	6.8	7.6	2.6	21.2	2.5				61.3			
Lateolabracidae	<i>Lateolabrax japonicus</i>	PM	0.5		0.3			1.3			0.3		2.3	5.6		4.7			6.6			6.6	23.4			
Haemulidae	<i>Plectorhinchus cinctus</i>	ZS					0.3			0.3			0.5				0.1			0.8			0.8			
Sparidae	<i>Acanthopagrus latus</i>	SM	1.0										1.0	1.3									1.3			
	<i>Acanthopagrus schlegelii</i>	SM	2.3	37.5	15.8	42.5	6.0	1.0	36.8		4.3		146.0	3.0	42.9	21.5	45.6	9.2	1.1	34.4		10.1	167.7			
	<i>Pagrus major</i>	SM		0.8		1.3		0.3	3.0		0.3	8.5	14.0		1.4		2.5		0.9	6.0		1.0	55.8	67.5		
Sillaginidae	<i>Sillago japonica</i>	SM				0.3	0.5	1.0	4.5	2.8	1.0	1.0	11.0				1.2	0.03	3.7	3.4	4.2	7.2	0.3	20.0		
Embiotocidae	<i>Ditrema viride</i>	ZS									0.5		0.5									7.0	7.0			

	<i>Ditrema temminckii pacificum</i>	ZS				4.5	1.0	0.5			0.3		6.3			82.3	15.0	8.2			4.8		110.3	
	<i>Ditrema</i> sp.	ZS								0.3			0.3								5.0		5.0	
	<i>Neoditrema ransonnetii</i>	ZS	2.3	1.3							0.5	4.0	19.8	12.1							5.9		37.8	
Labridae	<i>Parajulis poecilopterus</i>	ZS	0.3									0.3	8.1										8.1	
	<i>Halichoeres tenuispinis</i>	ZS					0.5					0.5					3.9						3.9	
Hexagrammidae	<i>Hexagrammos agrammus</i>	ZS	0.8		0.5	0.5	0.8				0.3	2.8	8.9		5.4	24.4	11.9				4.6		55.2	
	<i>Hexagrammos otakii</i>	ZS	1.3					0.3				1.5	8.0					3.6					11.6	
Cottidae	<i>Pseudoblennius cottoides</i>	ZS		0.3	3.3		3.8	0.3		0.3	0.3	8.0		1.2	11.0		23.5	1.4		0.9	0.9		38.8	
Blenniidae	<i>Petroscirtes breviceps</i>	ZS		0.5	0.5			0.3				1.3		0.5	0.5			0.1					1.1	
Callionymidae	<i>Repomucenus curvicornis</i>	SM							0.3			0.3							0.0				0.0	
	<i>Repomucenus ornatipinnis</i>	SM				0.3		0.3				0.5				2.9			0.0				2.9	
	<i>Repomucenus beniteguri</i>	SM									0.8	1.0	1.8							0.8	10.7		11.5	
Gobiidae	<i>Luciogobius guttatus</i>	SM				0.3						0.3				0.0							0.0	
	<i>Pterogobius elapoides</i>	ZS	1.8	1.0	0.3	0.5	0.5				0.3	4.3	9.4	6.3	2.0	0.2	4.3			1.4			23.6	
	<i>Tridentiger trigonocephalus</i>	SM		0.5							0.3	0.3	1.0		1.4					0.0	0.3		1.7	
	<i>Acentrogobius virgatus</i>	SM	4.3	8.5	3.3	0.3	0.8	8.8	0.5	2.8	9.8	0.8	39.5	9.5	14.5	9.2	0.4	1.4	11.6	0.3	3.3	11.1	1.0	62.3
	<i>Favonigobius gymnauchen</i>	SM	2.5	16.5	11.0	10.5	1.3	2.3	7.0	1.0	3.8	4.5	60.3	5.2	26.8	18.8	10.1	1.5	2.4	8.5	0.8	2.3	3.5	79.8
	<i>Gymnogobius heptacanthus</i>	ZS		0.3					0.3				0.5		0.2					0.0			0.2	
	<i>Chaenogobius gulosus</i>	ZS	1.5				0.3					1.8	1.6				0.2						1.7	
Siganidae	<i>Siganus fuscescens</i>	ZS			0.3					0.3		2.5	3.0			0.02				0.0		10.1	10.2	
Soleidae	<i>Zebrias zebrinus</i>	SM									0.3	0.3									0.1		0.1	
Monacanthidae	<i>Rudarius ercodes</i>	ZS	0.5	0.8	0.8	4.3	8.3	1.0	2.5	9.0	12.3	6.5	45.8	0.2	1.1	3.1	5.3	7.5	0.3	0.6	3.2	5.5	2.9	29.7
	<i>Thamnaconus modestus</i>	ZS	0.8	0.5		1.8	0.3	0.5					3.8	4.0	1.6		8.0	0.4	4.4				18.2	
	<i>Stephanolepis cirrifer</i>	ZS			1.0	1.8	0.3					6.3	9.3			4.4	3.8	0.1				16.9	25.3	
Tetraodontidae	<i>Takifugu pardalis</i>	SM	0.8				0.5		0.3				1.5	10.8				1.7		1.5			14.0	
	<i>Takifugu poecilonotus</i>	SM		0.3		0.5						0.3	1.0		0.2		4.7				0.7		5.6	
	<i>Takifugu niphobles</i>	SM	13.5	2.8		0.5	1.5	0.3				0.5	19.0	16.9	3.9		0.3	1.6	0.2		1.6		24.4	

Pleuronectidae	<i>Pleuronectes yokohamae</i>	SM										0.3	0.3									0.9	0.9
Total			81.3	116.8	51.0	86.3	39.0	226.0	56.5	17.0	34.8	47.8		356.5	292.6	138.2	332.9	130.8	518.3	58.8	18.5	58.6	152.1
WT (°C)			24.8	25.2	25.8	27.8	26.5	26.1	24.7	26.9	25.1	26.9											
Salinity			32.6	32.1	31.5	31.6	31.3	31.1	27.8	30.8	33.0	33.0											

For Review Only

Table 2. Results of linear model to examine possible effect of environmental conditions (seagrass shoot density, salinity and water temperature) on fish community. Fish was divided into three groups depending on their habitats or lifestyles: pelagic or migratory species (PM), sand or mud bottom-associated species (SM) and seagrass (*Z. marina*)- or substrate (rocky bottom including macrophytes)-associated species (ZS). Response variable was the mean biomass of each group. Initial explanatory variables were seagrass shoot density, salinity, and water temperature.

Group	Analysis of variance table (Type II tests)					Summary of model			
	Source	df	SS	F	P	Parameter	Estimate	SE	P
ZS	Error	7	19771			Intercept	-761.142	370.305	0.079
	Seagrass	1	66856	23.671	0.002	Seagrass	3.387	0.696	0.002
	Salinity	1	12080	4.277	0.077	Salinity	24.356	11.777	0.077
SM						Intercept	45.970	8.592	<0.000
PM	Error	6	177.585			Intercept	26.790	54.861	0.643
	Seagrass	1	44.537	1.505	0.266	Seagrass	0.088	0.07183	0.266
	Salinity	1	60.955	2.060	0.201	Salinity	1.759	1.22562	0.201
	Water Temperature	1	87.237	2.947	0.137	Water Temperature	-3.062	1.78369	0.137

Adjusted R-squared: ZS= 0.75, SM= -0.46, PM= 0.28